

Package: drought (via r-universe)

October 16, 2024

Type Package

Title Statistical Modeling and Assessment of Drought

Version 1.2

Date 2024-03-19

Author Zengchao Hao

Maintainer Zengchao Hao <z.hao4univ@gmail.com>

Description Provide tools for drought monitoring based on univariate and multivariate drought indicators. Statistical drought prediction based on Ensemble Streamflow Prediction (ESP), drought risk assessments, and drought propagation are also provided. Please see Hao Zengchao et al. (2017) <[doi:10.1016/j.envsoft.2017.02.008](https://doi.org/10.1016/j.envsoft.2017.02.008)>.

Depends R (>= 3.5.0)

Imports stats,copula,corrplot

Suggests MASS

License GPL-3

RoxygenNote 7.1.2

Encoding UTF-8

NeedsCompilation no

Date/Publication 2024-03-19 12:40:02 UTC

Repository <https://drought.r-universe.dev>

RemoteUrl <https://github.com/cran/drought>

RemoteRef HEAD

RemoteSha 830baf16af6049716aa47a9dc2216bdfa6adb590

Contents

drought-package	2
ACCU	3
BiEmp	4

ESPPred	4
JDSI	5
MSDI	6
PropagationMCC	7
RunDS	8
SDI	9
UMFreq	9
UniEmp	10
Index	11

drought-package	<i>Statistical Modeling and Assessment of Drought</i>
-----------------	---

Description

Provide tools for drought monitoring based on univariate and multivariate drought indicators. Statistical drought prediction based on Ensemble Streamflow Prediction (ESP), drought risk assessments, and drought propagation are also provided. Please see Hao Zengchao et al. (2017) <doi:10.1016/j.envsoft.2017.02.008>.

Details

Package: drought
 Type: Package
 Version: 1.1
 License: GPL-3

References

- Hao, Z. et al. (2017), An integrated package for drought monitoring, prediction and analysis to aid drought modeling and assessment, *Environ Modell Softw*, 91, 199-209.
- Hao, Z., and V. P. Singh (2015), Drought characterization from a multivariate perspective: A review *J. Hydrol.*, 527
- Hao and AghaKouchak (2013) Multivariate Standardized Drought Index: A parametric multi-index model, *Advances in Water Resources* 57, 12-18.
- Kao, S. C. and R. S. Govindaraju (2010). A copula-based joint deficit index for droughts. *Journal of Hydrology*, 380(1-2): 121-134.
- Hao, Z. et al. (2014). Global integrated drought monitoring and prediction system. *Scientific Data*, 1

Examples

```

#' X=runif(120, min = 0, max = 100) # 10-year monthly data
#' Yc<-ACCU(X,ts=6) # Compute the 6 month accumulated series
#' fit1<-SDI(X,ts=6) # Get the standardized drought index (or SPI)

#' z=matrix(t(fit1$SDI),ncol=1)
#' Res <- RunDS(z, -1)# Get drought duration and severity based on threshold SPI=-1

#' Y=runif(120, min = 0, max = 100) # 10-year monthly data
#' fit2<-MSDI(X,Y,ts=6) # Compute the 6 month Multivariate Standardized Drought Index (MSDI)
#' fit2$MSDI #Get the empirical MSDI

#' PropagationMCC(X, Y, 12, c(-1,1)) # Plot drought propagation

```

ACCU

Obtain the accumulation of monthly hydro-climatic variables

Description

Obtain the accumulation of monthly hydro-climatic variables

Usage

```
ACCU(X, ts = 6)
```

Arguments

X	The vector of monthly hydro-climatic variables of n years. ts is the accumulated time scale.
ts	The accumulated time scale

Examples

```

X=runif(120, min = 0, max = 100) # 10-year monthly data
Y<-ACCU(X,ts=3) # Compute the 3 month accumulated series

```


Value

The prediction of univariate and multivariate drought index based on ESP

Examples

```
X=runif(120, min = 0, max = 100) # 10-year monthly data
Y=runif(120, min = 0, max = 100)
ESPPred(X,Y,L=1,m=7,ts=6)
```

JDSI

Compute Joint Drought Severity Index with joint distribution

Description

The JDSI can be computed based on joint distribution or kendall distribution

Usage

```
JDSI(X, Y, ts = 6, type = 1)
```

Arguments

X	is the vector of a monthly hydro-climatic variable of n years.
Y	is the vector of a monthly hydro-climatic variable of n years.
ts	is the accumulated time scale.
type	is the method used to compute the JDSI (1 is Joint distribution and 2 is the Kendall function).

Value

The multivariate drought index based on the joint distribution or Kendall distribution

References

Hao, Z. et al. (2017) An integrated package for drought monitoring, prediction and analysis to aid drought modeling and assessment, *Environ Modell Softw*, 91, 199-209.

Examples

```
X=runif(120, min = 0, max = 100) # 10-year monthly data
Y=runif(120, min = 0, max = 100) # 10-year monthly data
fit<-JDSI(X,Y,ts=6)
z=matrix(t(fit$JDSI),ncol=1)
plot(z, type="l", col=1, lwd=2, lty=1, xlim=c(0,120),xlab="Time",ylab="JDSI")
```

MSDI

Compute the Multivariate Standardized Drought Index (MSDI)

Description

Based on a pair of monthly hydro-climatic variable (or corresponding marginals), the MSDI is computed using the joint distribution (parametric or nonparametric forms). The current version is based on the Gringorten plotting position. It can be extended to higher dimensions, such as trivariate case including meteorological, agricultural, and hydrological droughts. For the high dimension case, the copula or vine copula method can be employed

Usage

```
MSDI(X, Y, ts = 6)
```

Arguments

X is the vector of a monthly hydro-climatic variable of n years.
Y is the vector of a monthly hydro-climatic variable of n years.
ts is the accumulated time scale.

Value

The monthly MSDI series of different time scales (based on Gringorten plotting position)

References

Hao and AghaKouchak (2013) Multivariate Standardized Drought Index: A parametric multi-index model, *Advances in Water Resources* 57, 12-18.

Examples

```
X=runif(120, min = 0, max = 100) # 10-year monthly data  
Y=runif(120, min = 0, max = 100) # 10-year monthly data  
fit<-MSDI(X,Y,ts=6) # Compute the 6 month drought index  
fit$ProbEmp2 #Get the empirical drought index (e.g.,Gringorten plotting position )
```

PropagationMCC

Compute drought propagation based on maximum correlation

Description

Compute the pearson correlation between multi-time scale SPI and 1-month SRI to reflect the most possible propagation time (PT) from meteorological drought to hydrological drought. Note here the propagation of meteorological to hydrological drought is used as an example. The propagation of other types of drought can also be computed.

Usage

```
PropagationMCC(X, Y, acc = 12, lim = c(-1, 1), color = NA)
```

Arguments

X	The vector of monthly meteorological variable (e.g., precipitation)
Y	The vector of monthly hydrological variables (e.g., runoff)
acc	Maximum of propagation time (or accumulation periods)
lim	The limits interval for color
color	Color vector in plot

Value

Plot of correlation matrix

References

Xu, Y. et al (2019). Propagation from meteorological drought to hydrological drought under the impact of human activities: A case study in northern China. *J. Hydrol.* 579, 124147. Zhang Y., Hao Z., Feng S., et al. (2021). Agricultural drought prediction in China based on drought propagation and large-scale drivers. *Agr. Water Manage.*, 255: 107028.

Examples

```
X=runif(120, min = 0, max = 100) # 10-year monthly data
Y=runif(120, min = 0, max = 100) # 10-year monthly data
acc <- 12
lim <- c(-1,1)
PropagationMCC(X, Y, acc, lim)
```

RunDS

Compute drought duration and severity based on run theory

Description

The input data is monthly drought indices. Duration is defined as the length of consecutive time series when drought index is below the threshold value (e.g., -1). Severity is defined as the summation of drought index below the threshold. This analysis based on run theory is also referred to as threshold level method. Here the standardized drought index (SDI) is used as the example to compute the drought characteristics. Other univariate and multivariate drought indices can also be used.

Usage

```
RunDS(DI, thre)
```

Arguments

DI	The vector of the drought index (e.g., monthly SPI)
thre	The threshold of drought index (e.g., -0.5,-1)

Value

The duration and severity of each drought event

References

Yevjevich V. (1967). An Objective Approach to Definitions and Investigations of Continental Hydrologic Droughts. Hydrology Paper 23. Colorado State University, Fort Collins, CO.

Examples

```
X=runif(120, min = 0, max = 100) # 10-year monthly data
thre=-1 # specify the threshold value
fit<-SDI(X,ts=3) # Compute the univariate drought index, such as SPI
z=matrix(t(fit$SDI),ncol=1) # Reshape the matrix to a vector
Res <- RunDS(z, thre) # Compute the duration and severity
```

SDI *Compute the standardized drought index*

Description

Based on the vector of monthly variables, the standardized drought index is computed. Note here the standardized precipitation index (SPI) is used as the example of the drought index in the univariate case. It also represents other drought indices computed in the similar way as SPI.

Usage

```
SDI(X, ts = 6, dist = "EmpGrin")
```

Arguments

X The vector of a monthly hydro-climatic variable of n years.
ts is the accumulated time scale.
dist is a distribution function. The inputs can be "EmpGrin", "EmpWeib", "Gamma", "Lognormal".

Details

Apart from the standardized drought index, the percentile (probability) is also provided,

Value

The (univariate) standardized drought index of different time scales from both the empirical and parametric distribution

Examples

```
X=runif(120, min = 0, max = 100) # 10-year monthly data
fit<-SDI(X,ts=3) # Compute the 3 month drought index
fit$SDI # Get the empirical drought index
z=matrix(t(fit$SDI),ncol=1)
plot(z, type="l", col=1, lwd=2, lty=1, xlim=c(0,120),xlab="Time",ylab="SDI")
```

UMFreq *Univariate and multivariate return period*

Description

Univariate and multivariate return period

Usage

```
UMFreq(X, Y, EL = 1)
```

Arguments

X is the drought properties (e.g., duration) or indices (e.g., SPI)
Y is the drought properties (e.g., duration) or indices (e.g., SRI)
EL is the average recurrence time

Value

The univariate and multivariate return period

Examples

```
X=runif(60, min = 0, max = 100) # 60 drought duration values or index values  
Y=runif(60, min = 0, max = 100)  
fit<-UMFreq(X,Y,1)
```

UniEmp

Compute the univariate empirical joint probability (EMP)

Description

Compute the univariate empirical joint probability (EMP)

Usage

```
UniEmp(X, dist = "Gringorten")
```

Arguments

X The vector of a monthly hydro-climatic variable of n years.
dist is the function for the plotting position formula (Gringorten or Weibull).

Value

The univariate EMP

Examples

```
X=runif(20, min = 0, max = 100) # 20 monthly values of precipitation (e.g., August)  
fit<-UniEmp(X,dist = "Gringorten")
```

Index

ACCU, [3](#)

BiEmp, [4](#)

drought (drought-package), [2](#)

drought-package, [2](#)

ESPPred, [4](#)

JDSI, [5](#)

MSDI, [6](#)

PropagationMCC, [7](#)

RunDS, [8](#)

SDI, [9](#)

UMFreq, [9](#)

UniEmp, [10](#)